

# The White Spot Syndrome Virus: Emerging Savior or killer of Crustaceans' Future Generations?

Elida Cleyse Gomes da Mata Kanzaki<sup>1</sup>, Isamu Kanzaki<sup>1\*</sup>

<sup>1</sup>Laboratory of Bioprospection, University of Brasilia, DF, Brazil. CEP 70910-900.

## ABSTRACT

Emerging viruses pose enormous challenge to humanity. The sole member of the *Nimaviridae* family, in the genus *whispovirus*, the white spot syndrome virus, has spread to all around the world, and its infection cycle devastated shrimp aquaculture, menacing significantly the world market of food production. The comprehension of the mechanisms and factors involved in the emergence of novel viruses allow us to prevent and control the surge of new virus, and consequentially ensure our health and well-being, wherever avoiding human or animal disease acquisition, and particularly, to preserve food production from shrimp farming.

**Key words:** Whispovirus, shrimp, environment, endogenous virus.

## INTRODUCTION

Nutrient production is an essential activity of modern civilization that aims to constantly maintain food security, as defined by the World Food Summit in 1996 (1). Marine aquaculture systems have an important role in the global food supply chain (2), and demands extraordinary care to be protected from a variety of adverse factors, including infection and diseases by harmful microbial agents, mainly expressed in confined and altered environments as it happens in farming of aquatic metazoa organisms (3). Particularly referring to shrimp farming, a viral disease erupted in 1992-1993, in Asia, devastating marine shrimp production in captivity (4,5), rapidly expanding to other continents, imposing surmount economic losses in various countries dedicated to high density shrimp farming (5,6).

According to Santos et al. (2013), a crustaceans' disease etiologically linked to the White Spot Syndrome Virus (WSSV) has impacted the global economy, mainly on shrimp farming, with a loss of one billion dollar by year (7). In Brazil, the initial reports of the WSSV infection of farming shrimps, *Litopenaeus vanamei*, was reported in 2005, causing huge impact in the local economy, in the Laguna area of Santa Catarina State, with shrimp mortality reaching 80 %, and

loosing approximately 3 million dollars at that time. Also, other Brazilian regions, in the coastal northeast, representing the most productive areas of shrimps in captivity, were also heavily affected (8).

The etiological agent of the White Spot Syndrome was initially misunderstood as a baculovirus, as it is encompassed in the group of "the nuclear arthropod large DNA viruses" (9-11). The WSSV is presently classified by the International Committee on Taxonomy of Viruses as a member of the *Nimarividae* family, genus *Whispovirus*, considered to be the largest virion infecting the animal phyla (11). The virus isolation was firstly reported by Inouye et al. (1994) from, kuruma shrimp, *Penaeus japonicus*, in 1993, from farm ponds in western Japan (4), even though, the characteristic disease was previously reported from shrimp farms in Taiwan and Korea (12,13).

WSSV infected shrimps develop 2-6 mm white spots in the inner side of the cuticle, and body's red discoloration. Clinically, in experimental conditions, the disease can be asymptomatic, progressing to the anorexic state and the final death, that usually takes place 36 hours post-infection (4,12,13). The main pathological features of the white spot syndrome in *Penaeus japonicus* refer to epithelial cells'

### Address for Correspondence:

Isamu Kanzaki, Laboratory of Bioprospection, University of Brasilia, DF, Brazil. CEP 70910-900.

DOI: <https://doi.org/10.33309/2639-8893.060101>

© 2025 The Author(s). This open access article is distributed under a Creative Commons Attribution (CC-BY) 4.0 license.

degenerative lesions of the cuticular epidermis and stomach, as also of the connective tissues and lymphoid organs (11,13). Virions display an outer envelope lipoprotein, and a protein tegument linking the envelope to the nucleocapsid, exhibiting a robust and long oval shape morphology, measuring  $353 \pm 55$  nm and  $179 \pm 22$  nm, by cryogenic electron microscopy technique. Viral particles were described with and without tail-like structures. The nucleocapsid, assembled by 14 capsomere units, generates a ring-like structure proposed to conform a helix or stacked-ring structure linked by filaments, that in any case, packs the double circular DNA strain of 305-307 kbps (14,15).

In the infectious cycle, the *per os* infectivity factor (PIF) complex plays a major role in the initial process of virion internalization in the host cell (11,16). Either way, the viral envelope protein 28 (VP28) attaches to the host cells' membrane proteins, and complemented by the interaction of the virion tegument protein 24 (VP24) to multiple cell receptors, among them, the transmembrane polymeric immunoglobulin receptor (pIgR) as also the  $\text{Na}^+\text{-K}^+$ -ATPase alpha subunit (PvATP1A), the virion envelope fuses to the cytoplasmic cell membrane. The pIgR and PvATP1A are extensively expressed in shrimp cells, allowing WSSV infection of the epidermis, gills, foregut, hindgut, lymphoid organ, muscle, heart, and gonads. The binding of VP24 to the pIgR or PvATP1A promotes the clathrin-dependent or the caveolin-mediated virion endocytosis and/or micropinocytosis associated to the AP-2 protein (17,18). The endocytosed virion is coupled to the cell's calmodulin which triggers the mTORC1 signaling pathway, ultimately promoting viral protein translation (19). Previously, the viral envelope is merged to the endosome membrane, to traffic the capsid to the nucleopore, mediated by cellular microtubules, and transported by CgImportin  $\alpha 1/\beta 1$  to the nuclear compartment (20), where the viral DNA is driven-pressure ejected from the viral capsid (21). The WSSV genome is composed of more than 180 open reading frames (ORFs), most of them coding at least for 59 structural proteins, of them, 35 are assembled in the envelope and tegument, and 9 proteins constitute the nucleocapsid (22,23). Despite these early findings, the complete genome of a WSSV Korean isolate yielded 515 ORFs, among them 90 ORFs had any homology to known proteins in the current databases. The other 425 ORFs correspond to translated products of enzymes participating in the metabolism of nucleotides, genome replication and transcription besides main structural proteins (23). Marks et al. (2006) experiments results suggest that the viral DNA immediate early genes' transcription operates via the host cell RNA polymerase II transcription machinery (24). Non-structural proteins transcribed from the immediate early genes are mostly transcription factors, essential for viral DNA replication, which modulates the expression of both viral and host cell genes. Early and late genes, regulated by the immediate early gene products, express structural proteins, which are transported to the cell's nuclear compartment for virion assembly (25). The opposite poles of the oval form

nucleocapsids exhibit a portal cap and a closed base. The portal cap works as a complex molecular motor that translocate the viral genome into the capsid, whilst the closed base is open during DNA ejection, which suggested by electron microscopy, an image of filaments at this capsid pole, of both ejected DNA and DNA associated proteins. Mature capsids, containing the DNA genome, are surrounded by the tegument and envelope. All processes of virion morphogenesis take place in the host cell nuclear compartment, and the virion progeny are released by cell lysis (14,20).

Naturally, crustaceans display a diversity of antiviral mechanisms. Commonly, host cell pattern recognition receptors recognize the WSSV through their expressed pathogen associated molecular patterns leading to the activation of the crustacean immune system, the cellular and humoral response, eventually activating the transcription of antimicrobial genes. So, innumerable antimicrobial peptides exert antiviral activity by intercalating themselves in the viral envelope, which disturb the integrity and function of the envelope viral proteins involved in the attachment and interaction with host cell membrane proteins and virus cell receptors, as also other mechanisms during the stages of virus cell infection (26-29). Other host cell mechanisms look for defense against viral infection as the RNA interference (30-33), the circular RNA named the Quaking RNA-binding protein (34,35), or Heat Shock Protein 70 (36), the C3 like complement factor (37), among others. Nevertheless, virus encoded factors counteract host cell immune mechanisms or utilize the antiviral factors for its own benefit to accomplish the infection process (38).

## DISCUSSION

The emergence of new viruses is subjected to many factors considering that the hijacked host cell biochemical machinery is utilized by the virus molecules to guide the synthesis of its own molecules including the genome replication and transcription to yield the viral structural and non-structural proteins, including key enzymes, as also necessary molecular modifications to assemble new progeny virions (14-27). During this process, host cell via different pathways counter acts the viral infection mechanism (28-37). Successful virus infection and progeny release from host cell is usually accompanied by virus genome mutations caused by the cell's selective pressure, and also inherent errors of virus and cell's polymerases mechanisms to generate virus genome and transcripts, as also the failure of the enzyme's repair mechanisms (38,39).

Besides the mechanisms of host cell virus interaction, environment factors play a significative role in the evolution of viruses (40). Crustacean's production in captivity involves different scenarios, as the density of cultivated species, quality of water determined by animal's excreta as the pH, salinity, turbidity and so on. All these external pressure factors demand crustaceans' adaptation to the new environment, which could

lead to the selection of adapted ones, and death of unfit animals (41-43).

Some researchers have found possible WSSV ancestrals, such as endogenous nimaviruses integrated in different species of crustaceans. One of these endogenous viral elements were found in the genome of the marine white leg shrimp, *Penaeus vannamei*. So, 43 WSSV genes sequences are closely related to the endogenous nimavirus, which encodes the dUTPase and nucleocapsid proteins (44). Other work detected an endogenous nimavirus in the Jamaican bromeliad crab, *Metopaulias depressus*, sharing 68 putative ORFs with the WSSV genome, that code for non-structural proteins participating in the mechanisms of replication and nucleotide metabolism, and also code for structural proteins of the nucleocapsid, tegument and envelope (45). Also, from the genome of the isopod crustacean *Armadillidium vulgare*, gene fragments coding for peptides of 42 and 43 aminoacids were detected with 73 % to 74 % homology to peptides of the WSSV envelope (46,47). The search in the crustacean genome database, of 14 species, revealed five novel nimavirus genomes, encompassing 28 core genes, including proteins of unknown function, homologs of 5 baculovirus PIF and a sulfhydryl oxidase which denotes that nimavirus and baculovirus share the same phylogenetic origin (48). Most recently, Hirono's group proposed the existence 2 major WSSV phylotypes that time diverged, estimated by the Bayesian statistical analysis, between 1970 and early 1980, concluding that the phylotype I and the ancestor nimavirus were etiologically involved in the 1990s pandemic, and the phylotype II was restricted to Asia and Australia. In addition, 2 cross-phylotype recombinants have been detected (49).

## CONCLUSION

Based on the hypothesis that host cells, under environmental stress, work as epigenetic factors (3,50,51) that wake-up endogenous genic segments, and promote the genomic organization of these endogenous elements, that are capable to assembly in mature viral particles, enabling the new progeny virions to leave the host cell in a "instinct" to survive and to keep essential genes for future generations (52). Therefore, the novel exogenous virus continues the exercise of infecting new host cells to maintain the existence of "running genes", now as viral genes. It is interesting to note that since the emergence of WSSV, the mortality in the elapsed time was fast and very high, reaching 100 % (4,5,13,14), besides the initially very virulent isolates of WSSV had bigger genome than the less virulent and recent ones (53-55), possibly explained by the deletion of unnecessary genes which would optimize the virus' mechanisms in its infection cycle. Certainly, the host cell and WSSV interactions will, time by time, attenuate the virus pathogenicity. Let's say in a colloquial form, "the rude virus is educated by the cell biochemical machinery to behave", instead of a "disastrous virus that destroys of all host cells". It becomes more "cautious and intelligent", it replicates, producing millions of new progeny virus and

dispose of host cell to maintain its genes, until the time it is safe to endogenize itself and keep the genes useful for the host organism. All biochemical tools for all processes of genome organization, virion assembly and budding or cell lysis as also, the mechanisms of viral genome endogenization, are available by the host cell molecular paraphernalia (56-58).

Both RNA and DNA viruses have endogenous relatives (58,59). There are evidences suggesting that inhospitable cell's microenvironment triggers virus evasion, and it would explain the emergence of new viruses and catastrophic epidemics (48,49,60).

## REFERENCES

1. FAO: Rome Declaration on World Food Security and World Food Summit Plan of Action. World Food Summit. Rome, Italy, FAO, November 13–17, 1996.
2. Falconer L, Cutajar K, Krupandan A, Capuzzo E, Corner RA, Ellis T, Jeffery K, Mikkelsen E, Moore H, O'Beirn FX, O'Donohoe P, Ruane NM, Shilland R, Tett P, Telfer TC. Planning and licensing for marine aquaculture. *Rev Aquac.* 2023;15(4):1374-1404. doi: 10.1111/raq.12783.
3. Wikumpriya GC, Prabhatha MWS, Lee J, Kim CH. Epigenetic Modulations for Prevention of Infectious Diseases in Shrimp Aquaculture. *Genes (Basel).* 2023;14(9):1682. doi: 10.3390/genes14091682.
4. Inouye K, Miwa S, Oseko N, Nakano H, Kimura T, Momoyama K, Hiraoka M. Mass mortalities of cultured kuruma shrimp *Penaeus japonicus* in Japan in 1993: electron microscopic evidence of the causative virus. *Gyobyo Kenkyu (Fish Pathol)*, 1994; 29, 149-158.
5. Cox N, De Swaef E, Corteel M, Van Den Broeck W, Bossier P, Nauwynck HJ, Dantas-Lima JJ. Experimental Infection Models and Their Usefulness for White Spot Syndrome Virus (WSSV) Research in Shrimp. *Viruses.* 2024;16(5):813. doi: 10.3390/v16050813.
6. Kibenge FS. Emerging viruses in aquaculture. *Curr Opin Virol.* 2019;34:97-103. doi: 10.1016/j.coviro.2018.12.008.
7. Dos Santos R, Varela APM, Cibulski S, Lima F, Spilki F, Heinzelmann L, Luz R, Abreu P, Rohe P, Cavalli L. A brief history of white spot syndrome virus and its epidemiology in Brazil. *Virus Reviews and Research.* 2013;18. 1-. 10.17525/vrr.v18i1-2.63.
8. Neves SR de A, Martins PCC. Surgimento das doenças virais na Carcinicultura Brasileira: impactos e estratégias da gestão de saúde / Emergence of viral diseases in Brazilian shrimp farming: impacts and strategies health management. *Braz J Develop.* 2021;7(6):61925-44.
9. Chen X, Chen P, Wu D, Huang H, Chi X. A new baculovirus of cultured shrimps. *Sci China C Life Sci.* 1997;40(6):630-5. doi: 10.1007/BF02882693.
10. Nadala EC Jr, Tapay LM, Loh PC. Characterization of a non-occluded baculovirus-like agent pathogenic to penaeid shrimp. *Dis Aquat Organ.* 1998;33(3):221-9. doi: 10.3354/dao033221.

11. van Oers MM, Herniou EA, Jehle JA, Krell PJ, Abd-Alla AMM, Ribeiro BM, Theilmann DA, Hu Z, Harrison RL. Correction to: Developments in the classification and nomenclature of arthropod-infecting large DNA viruses that contain pif genes. *Arch Virol.* 2023;168(10):255. doi: 10.1007/s00705-023-05857-9. Erratum for: *Arch Virol.* 2023;168(7):182. doi: 10.1007/s00705-023-05793-8.
12. Wang HC, Hirono I, Maningas MBB, Somboonwiwat K, Stentiford G, Ictv Report Consortium. ICTV Virus Taxonomy Profile: Nimaviridae. *J Gen Virol.* 2019;100(7):1053-1054. doi: 10.1099/jgv.0.001248.
13. Park JH, Lee YS, Lee S, Lee Y. An infectious viral disease of penaeid shrimp newly found in Korea. *Dis Aquat Organ.* 1998;34(1):71-5. doi: 10.3354/dao034071.
14. Cox N, De Swaef E, Corteel M, Van Den Broeck W, Bossier P, Dantas-Lima JJ, Nauwynck HJ. The Way of Water: Unravelling White Spot Syndrome Virus (WSSV) Transmission Dynamics in *Litopenaeus vannamei* Shrimp. *Viruses.* 2023;15(9):1824. doi: 10.3390/v15091824.
15. Huang HJ, Tang SL, Chang YC, Wang HC, Ng TH, Garmann RF, Chen YW, Huang JY, Kumar R, Chang SH, Wu SR, Chao CY, Matoba K, Kenji I, Gelbart WM, Ko TP, Wang HA, Lo CF, Chen LL, Wang HC. Multiple Nucleocapsid Structural Forms of Shrimp White Spot Syndrome Virus Suggests a Novel Viral Morphogenetic Pathway. *Int J Mol Sci.* 2023;24(8):7525. doi: 10.3390/ijms24087525.
16. Li L, Hong Y, Huo D, Cai P. Ultrastructure analysis of white spot syndrome virus (WSSV). *Arch Virol.* 2020;165(2):407-412. doi: 10.1007/s00705-019-04482-9.
17. Wang X, Chen C, Zhang N, Chen Q, Zhang F, Liu X, Li F, Shi ZL, Vlask JM, Wang M, Hu Z. Functional Peroral Infectivity Complex of White Spot Syndrome Virus of Shrimp. *J Virol.* 2022;96(24):e0117322. doi: 10.1128/jvi.01173-22.
18. Niu GJ, Wang S, Xu JD, Yang MC, Sun JJ, He ZH, Zhao XF, Wang JX. The polymeric immunoglobulin receptor-like protein from *Marsupenaeus japonicus* is a receptor for white spot syndrome virus infection. *PLoS Pathog.* 2019;15(2):e1007558. doi: 10.1371/journal.ppat.1007558.
19. Zhou J, Zhang H, Wu G, Zhang Y, Aweya JJ, Tayyab M, Zhu J, Zhang Y, Yao D. The Na<sup>+</sup>-K<sup>+</sup>-ATPase alpha subunit is an entry receptor for white spot syndrome virus. *mBio.* 2025;16(3):e0378724. doi: 10.1128/mbio.03787-24.
20. Hong PP, Li C, Niu GJ, Zhao XF, Wang JX. White spot syndrome virus directly activates mTORC1 signaling to facilitate its replication via polymeric immunoglobulin receptor-mediated infection in shrimp. *PLoS Pathog.* 2022;18(9):e1010808. doi: 10.1371/journal.ppat.1010808.
21. Li DL, Yang MH, Liu LK, Meng C, Li MQ, Liu HP. Invasion and Propagation of White Spot Syndrome Virus: Hijacking of the Cytoskeleton, Intracellular Transport Machinery, and Nuclear Import Transporters. *J Virol.* 2022;96(12):e0220521. doi: 10.1128/jvi.02205-21.
22. Sun M, Liu M, Shan H, Li K, Wang P, Guo H, Zhao Y, Wang R, Tao Y, Yang L, Zhang Y, Su X, Liu Y, Li C, Lin J, Chen XL, Zhang YZ, Shen QT. Ring-stacked capsids of white spot syndrome virus and structural transitions with genome ejection. *Sci Adv.* 2023;9(8):eadd2796. doi: 10.1126/sciadv.add2796.
23. Tan Yw, Shi ZL. Proteomic analyses of the shrimp white spot syndrome virus. *Virol Sin.* 2008; 23, 157–166. <https://doi.org/10.1007/s12250-008-2924-0>.
24. Marks H, Vorst O, van Houwelingen AMML, van Hulten MCW, Vlask JM. Gene-expression profiling of White spot syndrome virus in vivo. *J Gen Virol.* 2005;86(Pt 7):2081-2100. doi: 10.1099/vir.0.80895-0.
25. Chai CY, Yoon J, Lee YS, Kim YB, Choi TJ. Analysis of the complete nucleotide sequence of a white spot syndrome virus isolated from Pacific white shrimp. *J Microbiol.* 2013;51(5):695-9. doi: 10.1007/s12275-013-3171-0.
26. Marks H, Ren XY, Sandbrink H, van Hulten MC, Vlask JM. In silico identification of putative promoter motifs of White Spot Syndrome Virus. *BMC Bioinformatics.* 2006;7:309. doi: 10.1186/1471-2105-7-309.
27. Chen Y, Wu G, Wang C, Zhang H, Zhu J, Zhang Y, Lin Z, Yao D. Comparative Transcriptome Analysis Reveals That WSSV IE1 Protein Plays a Crucial Role in DNA Replication Control. *Int J Mol Sci.* 2022;23(15):8176. doi: 10.3390/ijms23158176.
28. Nupan B, Phongdara A, Saengsakda M, Leu JH, Lo CF. Shrimp Pm-fortilin inhibits the expression of early and late genes of white spot syndrome virus (WSSV) in an insect cell model. *Dev Comp Immunol.* 2011;35(4):469-75. doi: 10.1016/j.dci.2010.11.016.
29. Shashikumar A, Desai PV. Susceptibility of testicular cell cultures of crab, *Scylla serrata* (Forsk.) to white spot syndrome virus. *Cytotechnology.* 2013;65(2):253-62. doi: 10.1007/s10616-012-9482-x.
30. Liu LK, Liu MJ, Li DL, Liu HP. Recent insights into anti-WSSV immunity in crayfish. *Dev Comp Immunol.* 2021;116:103947. doi: 10.1016/j.dci.2020.103947.
31. Bin Hafeez A, Jiang X, Bergen PJ, Zhu Y. Antimicrobial Peptides: An Update on Classifications and Databases. *Int J Mol Sci.* 2021;22(21):11691. doi: 10.3390/ijms222111691.
32. Urmi UL, Vijay AK, Kuppusamy R, Islam S, Willcox MDP. A review of the antiviral activity of cationic antimicrobial peptides. *Peptides.* 2023;166:171024. doi: 10.1016/j.peptides.2023.171024.
33. Hong SJ, Kim KH. RNA interference targeting WSSV ribonucleotide reductase 2 provides long-term protection against infection in *Litopenaeus vannamei*. *Dis Aquat Organ.* 2024;159:71-78. doi: 10.3354/dao03805.
34. Limkul S, Phiwthong T, Wanvimonsuk S, Seabkongseng T, Aunkam P, Jaree P, Luangtrakul W, Mahanil K, Teamtisong K, Tittabutr P, Teamroong N, Sarnow P, Wang HC, Somboonwiwat K, Boonchuen P. Viral circular RNA-encoded protein, ceVP28, divulges an antiviral response in invertebrates. *Proc Natl Acad Sci U S A.* 2025;122(8):e2321707122. doi: 10.1073/pnas.2321707122.
35. Phiwthong T, Limkul S, Aunkam P, Seabkongseng T, Teamroong N, Tittabutr P, Boonchuen P. Quaking RNA-Binding protein (QKI) mediates circular RNA biogenesis in *Litopenaeus vannamei* during WSSV infection. *Fish Shellfish Immunol.* 2025;159:110178. doi: 10.1016/j.fsi.2025.110178.



36. Janewanthanakul S, Supungul P, Tang S, Tassanakajon A. Heat shock protein 70 from *Litopenaeus vannamei* (LvHSP70) is involved in the innate immune response against white spot syndrome virus (WSSV) infection. *Dev Comp Immunol*. 2020;102:103476. doi: 10.1016/j.dci.2019.103476.
37. Liu Y, Song Q, Li D, Zou R, Zhang Y, Hao S, Geng X, Sun J. A novel complement C3 like gene (Lv-C3L) from *Litopenaeus vannamei* with bacteriolytic and hemolytic activities and its role in antiviral immune response. *Fish Shellfish Immunol*. 2019;91:376-387. doi: 10.1016/j.fsi.2019.05.045.
38. Li C, Weng S, He J. WSSV-host interaction: Host response and immune evasion. *Fish Shellfish Immunol*. 2019;84:558-571. doi: 10.1016/j.fsi.2018.10.043.
39. Hill T, Unckless RL. Recurrent evolution of high virulence in isolated populations of a DNA virus. *Elife*. 2020;9:e58931. doi: 10.7554/eLife.58931.
40. Jones JE, Le Sage V, Lakdawala SS. Viral and host heterogeneity and their effects on the viral life cycle. *Nat Rev Microbiol*. 2021;19(4):272-282. doi: 10.1038/s41579-020-00449-9.
41. Poss M. Virus dynamics and evolution: bridging scales and disciplines. *Viruses*. 2011;3(8):1432-8. doi: 10.3390/v3081432.
42. Peng SE, Lo CF, Lin SC, Chen LL, Chang YS, Liu KF, Su MS, Ko GH. Performance of WSSV-infected and WSSV-negative *Penaeus monodon* postlarvae in culture ponds. *Dis Aquat Organ*. 2001;46(3):165-72. doi: 10.3354/dao046165.
43. Min JG, Kim YC, Kim KI. Role of Filter-Feeding Bivalves in the Bioaccumulation and Transmission of White Spot Syndrome Virus (WSSV) in Shrimp Aquaculture Systems. *Pathogens*. 2024;13(12):1103. doi: 10.3390/pathogens13121103.
44. Bao W, Tang KFJ, Alcivar-Warren A. The Complete Genome of an Endogenous Nimavirus (*Nimav-1\_LVa*) From the Pacific Whiteleg Shrimp *Penaeus (Litopenaeus) Vannamei*. *Genes (Basel)*. 2020;11(1):94. doi: 10.3390/genes11010094.
45. Rozenberg A, Brand P, Rivera N, Leese F, Schubart CD. Characterization of fossilized relatives of the White Spot Syndrome Virus in genomes of decapod crustaceans. *BMC Evol Biol*. 2015;15:142. doi: 10.1186/s12862-015-0380-7.
46. Thézé J, Leclercq S, Moumen B, Cordaux R, Gilbert C. Remarkable diversity of endogenous viruses in a crustacean genome. *Genome Biol Evol*. 2014;6(8):2129-40. doi: 10.1093/gbe/evul63.
47. Li H, Zhu Y, Xie X, Yang F. Identification of a novel envelope protein (VP187) gene from shrimp white spot syndrome virus. *Virus Res*. 2006;115(1):76-84. doi: 10.1016/j.virusres.2005.07.007.
48. Kawato S, Shitara A, Wang Y, Nozaki R, Kondo H, Hirono I. Crustacean Genome Exploration Reveals the Evolutionary Origin of White Spot Syndrome Virus. *J Virol*. 2019;93(3):e01144-18. doi: 10.1128/JVI.01144-18.
49. Kawato S, Omine R, Teruya S, Kubo H, Yasumoto S, Kondo M, Takahashi Y, Nozaki R, Kondo H, Hirono I. Evolutionary genomics of white spot syndrome virus. *Fish Sci*. 2023; 89: 769–783. <https://doi.org/10.1007/s12562-023-01715-4>.
50. Perera BPU, Silvestre F. Environmental Epigenomes. *Epigenomes*. 2023;7(3):21. doi: 10.3390/epigenomes7030021.
51. Zhang X, Hu C, Chen T, Li P, Tan Y, Ren C, Wang Y, Jiang X, Ma B, Yin J, Huang Y, Liu L, Li H, Luo P. DNA methylation regulates growth traits by influencing metabolic pathways in Pacific white shrimp (*Litopenaeus vannamei*). *BMC Genomics*. 2025;26(1):511. doi: 10.1186/s12864-025-11688-6.
52. Kanzaki LIB. HTLV-1: A real pathogen or a runaway guest of a diseased cell? *J Biosci*. 2018;43(4):785-795.
53. Li F, Gao M, Xu L, Yang F. Comparative genomic analysis of three white spot syndrome virus isolates of different virulence. *Virus Genes*. 2017;53(2):249-258. doi: 10.1007/s11262-016-1421-z.
54. Puttiringroj P, Kawato S, Mwamburi SM, Furukawa M, Oomine R, Koiwai K, Kondo H, Hirono I. Comparative genomics highlights the virulence and evolutionary trajectory of white spot syndrome virus. *J Gen Virol*. 2024;105(11). doi: 10.1099/jgv.0.002042.
55. Zwart MP, Dieu BT, Hemerik L, Vlak JM. Evolutionary trajectory of white spot syndrome virus (WSSV) genome shrinkage during spread in Asia. *PLoS One*. 2010 Oct 14;5(10):e13400. doi: 10.1371/journal.pone.0013400.
56. Kawato S, Nozaki R, Kondo H, Hirono I. Integrase-associated niche differentiation of endogenous large DNA viruses in crustaceans. *Microbiol Spectr*. 2024;12(1):e0055923. doi: 10.1128/spectrum.00559-23.
57. da Mata Kanzaki ECG, Kanzaki I. Viral Genome Integration into the Host Cell Genome: A Double Edged-Sword. *Discov Med*. 2021;32(167):141-148.
58. Déjosez M, Marin A, Hughes GM, Morales AE, Godoy-Parejo C, Gray JL, Qin Y, Singh AA, Xu H, Juste J, Ibáñez C, White KM, Rosales R, Francoeur NJ, Sebra RP, Alcock D, Volkert TL, Puechmaille SJ, Pastusiak A, Frost SDW, Hiller M, Young RA, Teeling EC, García-Sastre A, Zwaka TP. Bat pluripotent stem cells reveal unusual entanglement between host and viruses. *Cell*. 2023;186(5):957-974.e28. doi: 10.1016/j.cell.2023.01.011.
59. Moniruzzaman M, Aylward FO. Endogenous DNA viruses take center stage in eukaryotic genome evolution. *Proc Natl Acad Sci U S A*. 2023;120(21):e2305212120. doi: 10.1073/pnas.2305212120.
60. Kreiling JA. Dysregulation of endogenous retroviruses triggers aging and senescence. *Nat Aging*. 2024;4(12):1670-1672. doi: 10.1038/s43587-024-00759-0.

**How to cite this article:** Elida Cleyse Gomes da Mata Kanzaki, Isamu Kanzaki. The White Spot Syndrome Virus: emerging savior or killer of crustaceans' future generations?. *J Pathol Infect Dis* 2025;6(1): 1-5.

DOI: <https://doi.org/10.33309/2639-8893.060101>